

# Mathematics II – Examples

## II.7.\* Derivative of composed function

Let the functions  $z = f(x, y)$ ,  $x = x(u, v)$ ,  $y = y(u, v)$  are differentiable. Then

$$\frac{\partial z}{\partial u} = \frac{\partial f}{\partial x} \cdot \frac{\partial x}{\partial u} + \frac{\partial f}{\partial y} \cdot \frac{\partial y}{\partial u} \quad (1)$$

$$\frac{\partial z}{\partial v} = \frac{\partial f}{\partial x} \cdot \frac{\partial x}{\partial v} + \frac{\partial f}{\partial y} \cdot \frac{\partial y}{\partial v} \quad (2)$$

**Example 160:** For the following differentiable functions  $z = f(x, y)$ ,  $x = u \cos v$ ,  $y = u \sin v$

- Compute the derivative of composed function  $\frac{\partial z}{\partial u}$ ,  $\frac{\partial z}{\partial v}$ .
- Calculate for the particular function  $z = f(x, y) = e^x \ln y$ .

**Example 161:**  $w = f(x^4 + y^4 - 2z^4)$ . Compute  $V = \frac{1}{4x^3} \frac{\partial w}{\partial x} + \frac{1}{4y^3} \frac{\partial w}{\partial y} + \frac{1}{4z^3} \frac{\partial w}{\partial z}$ .

**Example 162:** Verify that the function  $y = f(x + at) + g(x - at)$  satisfies the partial differential equation  $\frac{\partial^2 y}{\partial t^2} - a^2 \frac{\partial^2 y}{\partial x^2} = 0$ , where  $a$  is a constant in  $\mathbb{R}$ . Assume that both  $f$  and  $g$  have continuous partial derivative of second order. [Hint:  $u = x + at$ ,  $v = x - at$ ,  $y = f(u) + g(v)$ ]

**Example 163:** Verify that the function  $z = f\left(\frac{y}{x}\right)$  satisfies the equation  $x \frac{\partial z}{\partial x} + y \frac{\partial z}{\partial y} = 0$ . Assume that  $f$  is a differentiable function.

**Example 164:** Let us have  $z = f(u, v)$ ,  $u = x^2 - y^2$ ,  $v = e^{xy}$ . Compute the following differential expression  $W = y \frac{\partial z}{\partial x} + x \frac{\partial z}{\partial y}$ , where  $f$  is a differentiable function.

**Example 165:** Let  $f(u, v) = x^y$ , where  $x = u^2 + v^2$ ,  $y = uv + v^2$ . Compute  $\frac{\partial f}{\partial u}$  and  $\frac{\partial f}{\partial v}$  in the point  $A$ , which has coordinations  $u = 1$ ,  $v = -1$ .

## II.9. Implicit functions

**Example 166:** Prove, that the equation  $x^3 + y^3 = 2x^2 + xy - 1$  defines unique implicit function  $y = f(x)$  in a neighbourhood of the point  $A = [1, 0]$ . Compute  $f'(1)$  and  $f''(1)$ .

**Example 167:** Write equations of both a tangent line  $t$  and a normal line  $n$  in the point  $A = [1, 1]$  to the curve, which is defined by implicit way by the equation  $F(x, y) \equiv x^3 y + y^3 x + x^2 y - 3 = 0$ .

**Example 168:** By the equation  $\ln \sqrt{x^2 + y^2} = \arctg \frac{y}{x}$  for  $x \neq 0$  is defined an implicit function  $y = f(x)$ . Compute  $y'$  and  $y''$ .

**Example 169:** Show, that the equation  $x^2 + 2xy + y^2 - 4x + 2y - 3 = 0$  define an implicit function  $y = f(x)$ , which graph is passing through the point  $A = [0, 1]$ . Check, if the function  $f$  is convex in some neighbourhood of the point  $x_0 = 0$ . Write an equation of a tangent line to the graph of this function in the point  $A$

**Example 170:** Prove, that the equation  $F(x, y, z) \equiv z^3 + 3x^2 z - 2xy = 0$  defines unique implicit function  $z = f(x, y)$  in a neighbourhood of the point  $A = [-1, -2, 1]$ . Compute  $\text{grad} f(-1, -2)$ .

**Example 171:** In a neighborhood of the point  $[2, -2, 1]$  is given implicit function  $z = f(x, y)$  by the equation  $\ln z + x^2 y z + 8 = 0$ . Compute directional derivative  $\frac{\partial z}{\partial \vec{s}}(A)$ , where  $\vec{s} = \overrightarrow{AB}$ ,  $A = [2, -2]$ ,  $B = [3, -3]$ .

**Example 172:** Write equations of both a tangent plain  $\tau$  and a normal line  $n$  to a surface  $z = z(x, y)$  defined implicitly by the equation  $F(x, y, z) \equiv x^2 + y^2 + z^2 - 25 = 0$  in the point  $A = [3, 0, 4]$ .

**Example 173:** Write an equation of a tangent plain to a surface defined implicitly by the equation  $F(x, y, z) \equiv x(y + z) + z^2 - 5 = 0$  parallel with the plain  $\rho: 3x - 3y + 6z = 2$ .  
[Note: There are two tangent plains satisfying the given condition.]